

NUCLEAR MODEL OF BINDING ALPHA-PARTICLES *

K.A. GRIDNEV, S.YU. TORILOV

Institute of Physics, St.Petersburg State University, 198904, Russia

V.G. KARTAVENKO

Joint Institute for Nuclear Research, Dubna, Moscow District, 141980, Russia

W. GREINER, D.K. GRIDNEV

J.W. Goethe University, Frankfurt/Main, D-60054, Germany

J. HAMILTON

Vanderbilt University, Nashville, TN 37235, USA

The model of binding alpha-particles in nuclei is suggested. It is shown good (with the accuracy of 1-2%) description of the experimental binding energies in light and medium nuclear systems. Our preliminary calculations show enhancement of the binding energy for super heavy nuclei with $Z \sim 120$.

We assume that atomic nuclei might be considered as a tight packing of alpha-particles. The binding energy could be expressed as a sum of energies coming from interactions between N_α alpha-particles and their self-energies.

$$E_B(N_\alpha) = E_B(\alpha)N_\alpha + E_C + V_{\alpha\alpha}(N_\alpha) \quad (1)$$

where $V_{\alpha\alpha}(N_\alpha) = \sum_{j>i}^{N_\alpha} V_{\alpha\alpha}(r_{ij})$, $E_B(\alpha)=28.296$ MeV is the binding energy of alpha-particle, and E_C is the Coulomb part of the potential energy. For the alpha-alpha interaction we use the Lennard-Jones (LJ) potential, which can be written as

$$V_{\alpha\alpha}(r) = V_0 \left[(\sigma/r)^{12} - 2(\sigma/r)^6 \right] \quad (2)$$

where r is the distance between centers of clusters, V_0 is the depth of the

*This work is partially supported by Deutsche Forschungsgemeinschaft (grant 436 RUS 113/24/0-4), Russian Foundation for Basic Research (grant 03-02-04021) and the Heisenberg-Landau Program (JINR)




Figure 1. Separation energy of a single alpha-particle versus the number of alpha-particles N_α . The alpha-particle sets and the differences of inter-alpha-particle bonds ($\Delta N = B_\alpha(N_\alpha + 1) - B_\alpha(N_\alpha)$, where $B_\alpha(N_\alpha)$ is the number of the bonds for nucleus with N_α alpha-particles.)

potential ($V_{\alpha\alpha}(\sigma) = -V_0$, $dV_{\alpha\alpha}/dr|_{r=\sigma} = 0$), and σ is the pair bonding length. The positions of alpha-particles are found from the condition of




Figure 2. The set of distances between alpha-particles in the nucleus versus the number of bonds between alpha-particles B_α . Separation energy of an alpha-particle as the function of the number of alpha-particles N_α .

the minimal total potential energy. The volume of the nucleus is set on a grid and the first alpha-particle is positioned in the center. The potential is calculated in all nodes of the grid and the next particle is positioned into the node with the minimum potential energy ($r = \sigma$). Then we take two

particles and calculate their potential and put to its minimum the third cluster. And so on and so forth. This simple algorithm gives possibility to cast the binding energy (1) in the following form¹

$$E_B(N_\alpha) = E_B(\alpha)N_\alpha + V_0B_\alpha + E'_C \quad (3)$$

where the effective geometrical factor B_α is the number of bonds between alpha particles, and all Coulomb terms are collected in E'_C . Assuming six equal nucleon bonds in each close-packed tetrahedral alpha particle, the unit bond energy will approximately equal to $V_0 \approx E_B(\alpha)/6 \approx 5$ MeV.

The cluster structure of nuclear system, and the selected *type of interaction* (LJ Eq. (2) results to enhanced stability of tight cluster configurations with *icosahedral symmetry* (see Figures). It is seen, in the upper part of Fig. 2, that the first set of distances between alpha-particles are nearly $\sigma \approx 3$ fm, and then there are stripes corresponding to non-bordering alpha-particles and *etc.* Gradually the stripes wash out because the tetrahedrons have to deform in order to constitute closed figures.

The cluster structure of nuclear system leads to (many-body) shell effects (see Figures), which are not reduced to the traditional nucleon single-particle ones. Such shell effects could be very important for heavy and superheavy nuclei, which stability is defined mainly by the shell effects.

Assuming that the main trend remains when adding neutrons, our calculations show enhancement of the binding energy for super heavy nuclei with $Z \sim 120$, which approximately corresponded to the region of completed icosahedral shell of 55 alpha-particles. One of us ² has predicted recently fullerene-like nuclei in this region. The details (exact binding energy and the Z value) depend on the details of the potential, but the main trends conserve, as seen from the lower part of Fig. 2, where results are presented for two different values of the potential width ($R_W = 2^{1/12}\sigma((\sqrt{2}+1)^{1/6} - (\sqrt{2}-1)^{1/6})$).

It turns out also, the structure of heavy even-even nuclei looks like the Fe nucleus coated with the Bose condensate of alpha-particles with additional neutrons playing the role of electrons in the covalent bond. A similar mechanism of alpha-particle formation analogous with formation of Cooper pairs has been considered in ³.

References

1. P.D. Norman, *Eur. J. Phys.* **14**, 36 (1993).
2. W.Greiner, *Prog. Theor. Phys. Suppl.* **146**, 84 (2002).
3. S. Koh, *Prog. Theor. Phys. Suppl.* **132**, 197 (1998).